



TURKISH - FEPS PHYSIOLOGY CONGRESS

September 3-7, 2011

Yeditepe University

Istanbul



Abstract Submission Deadline 20 May 2011



www.feps2011.org



address: Cumhuriyet Cad. No:173/2 34373 Harbiye, Istanbul-Turkey
phone: +90 212 368 47 95 fax: +90 212 230 64 25
e-mail: feps2011@vip.com.tr



[PC137]

EVALUATION OF COGNITIVE DEFICITS IN PATIENTS WITH PARKIN (PARK2) GENE MUTATIONS USING AUDITORY EVENT-RELATED POTENTIALS

Atila Uslu¹, Mehmet Ergen², Hasan Demirci³, Ebba Lohmann⁴, Hasmet Hanagasi⁴, Tamer Demiralp¹

¹Department of Physiology, Istanbul Faculty of Medicine, Istanbul University, 34093 Capa, Istanbul, Turkey

²Department of Physiology, School of Medicine, Acibadem University, 34384 Maltepe, Istanbul, Turkey

³Department of Neuroscience, Research Institute for Experimental Medicine, Istanbul University, 34093 Capa, Istanbul, Turkey

⁴Department of Neurology, Behavioral Neurology and Movement Disorders Unit, Istanbul Faculty of Medicine, Istanbul University, 34093 Capa, Istanbul, Turkey

Objective: Studies on cognitive deficits in patients with parkin (PARK2) gene mutation are rare and report more slight changes in terms of cognitive deficits compared with other Parkinson's disease (PD) patients, probably because of the early-onset and slower progress of the disease in this group. The aim of the present study is to investigate the cognition in patients with parkin mutation more in depth using cognitive electrophysiological measures.

Methods: The participants consisted of 25 healthy control volunteers, 65 non-demented PD patients. PD patients were divided into three groups as: PD patients with parkin mutation (PM-PD; n=15) and early onset (EO-PD; n=25) and late onset (LO-PD; n=25) idiopathic PD patients. Neuropsychological state of the subjects was evaluated by Mini-Mental State Examination (MMSE) and ERPs were recorded while the subjects performed an auditory oddball task.

Results: We found MMSE scores of the all PD groups were significantly lower than those of the control group ($p < 0.001$). P300 amplitudes in target-ERPs of the oddball test were significantly lower in all PD groups compared with the controls ($p < 0.001$). However, P200 amplitudes of all PD groups were significantly larger than those of the control group ($p = 0.042$). In addition, the latencies of both P200 and P300 potentials were significantly longer in all PD patients ($p = 0.015$ and $p < 0.001$, respectively). There was no significant difference among the PD groups.

Conclusions: Our results suggest that EO-PD patients with PM-PD show a similar delay in cognitive processing and a similar decrease in selective attention performance as in other PD groups.

Go

[Meeting Index](#)
[Search](#)
[APH Homepage](#)
[Online Journal](#)
[Privacy Policy](#)
[Home](#)

Acta Physiologica Congress

[Back](#)

Acta Physiologica 2011; Volume 203, Supplement 686

Joint Congress of FEPS and Turkish Society of Physiological Sciences

03/09/2011-07/09/2011

Istanbul, Turkey

EVALUATION OF COGNITIVE DEFICITS IN PATIENTS WITH PARKIN (PARK2) GENE MUTATIONS USING AUDITORY EVENT-RELATED POTENTIALS

Abstract number: PC137

Uslu¹ Atilla, Ergen² Mehmet, Demirci³ Hasan, Lohmann⁴ Ebba, Hanagasi⁴ Hasmet, Demiralp¹ Tamer

¹*Department of Physiology, Istanbul Faculty of Medicine, Istanbul University, 34093 Capa, Istanbul, Turkey*

²*Department of Physiology, School of Medicine, Acibadem University, 34384 Maltepe, Istanbul, Turkey*

³*Department of Neuroscience, Research Institute for Experimental Medicine, Istanbul University, 34093 Capa, Istanbul, Turkey*

⁴*Department of Neurology, Behavioral Neurology and Movement Disorders Unit, Istanbul Faculty of Medicine, Istanbul University, 34093 Capa, Istanbul, Turkey*

Objective:

Studies on cognitive deficits in patients with parkin (PARK2) gene mutation are rare and report more slight changes in terms of cognitive deficits compared with other Parkinson's disease (PD) patients, probably because of the early-onset and slower progress of the disease in this group. The aim of the present study is to investigate the cognition in patients with parkin mutation more in depth using cognitive electrophysiological measures.

Methods:

The participants consisted of 25 healthy control volunteers, 65 non-demented PD patients. PD patients were divided into three groups as: PD patients with parkin mutation (PM-PD; n=15) and early onset (EO-PD; n=25) and late onset (LO-PD; n=25) idiopathic PD patients. Neuropsychological state of the subjects was evaluated by Mini-Mental State Examination (MMSE) and ERPs were recorded while the subjects performed an auditory oddball task.

Results:

We found MMSE scores of the all PD groups were significantly lower than those of the control group ($p < 0.001$). P300 amplitudes in target-ERPs of the oddball test were significantly lower in all PD groups compared with the controls ($p < 0.001$). However, P200 amplitudes of all PD groups were significantly larger than those of the control group ($p = 0.042$). In addition, the latencies of both P200 and P300 potentials were significantly longer in all PD patients ($p = 0.015$ and $p < 0.001$, respectively). There was no significant difference among the PD groups.

Conclusions:

Our results suggest that EO-PD patients with PM-PD show a similar delay in cognitive processing and a similar decrease in selective attention performance as in other PD groups.

To cite this abstract, please use the following information:

***Acta Physiologica 2011*; Volume 203, Supplement 686 :PC137**