### **PP-28**

## BTK PROTEIN DETERMINATION BY FLOW CYTOMETRY IN PATIENTS WITH X-LA

## Suzan CINAR, Immunology, Istanbul University, DETAE, Istanbul

Yuk Yin NG, Genetics, Istanbul University, DETAE, Istanbul

Abdullah YILMAZ, Immunology, Istanbul University, DETAE, Istanbul

Sinem FIRTINA, Department of Genetics, Istanbul University, Institute of Experimental Medicine (DETAE), Istanbul

Esra ÖZEK, Pediatrics Diseases, Istanbul University, Cerrahpasa Faculty of Medicine, Istanbul Safa BARIS, Paediatrics, Division of Pediatric Allergy and Immunology, Marmara University, Marmara University School of Medicine, Istanbul

Işıl BARLAN, Paediatrics, Division of Pediatric Allergy and Immunology, Marmara University, Marmara University School of Medicine, Istanbul

Yıldız CAMCIOĞLU, Pediatrics Diseases, Istanbul University, Cerrahpasa Faculty of Medicine, Istanbul

Uğur ÖZBEK, Genetics, Istanbul University, DETAE, Istanbul

Gunnur DENIZ, Immunology, Istanbul University, DETAE, Istanbul

X linked agammaglobulinemia (X-LA), is caused as a result of mutations in the Bruton's tyrosine kinase (Btk) gene and characterized by the discontinuation in stages of maturing and differentiation of B lymphocytes. Eightyfive % of agammaglobulinemia (AG) cases have mutations in Btk gene rendering in absence of mature B and plasma cells and very low levels of serum immunoglobulin. After the demonstration of absence of B cell and BTK expression with immunophenotyping by flow cytometry (FCM) in cases of AG, Btk gene mutation analysis can be carried out for the molecular genetic diagnosis of X-LA. The aim of this study is to determinate BTK expression in monocytes of AG patients and their mothers by FCM before BTK gene mutation test with PCR. We analysed 11 AG male patients having low Ig levels and CD19 expression with recurrent infections. In order to prove B cell deficiency, B cell expression has been determined by FCM using anti-CD19 monoclonal antibodies (mAb). Btk expression in B cells and monocytes were determined by using intracytoplasmic staining method with anti-BTK mAb after surface staining with anti-CD45 and -CD14 (Figure 1). Following this, DNA samples of the cases were screened for Btk gene mutation with PCR (Table 1). Five of 11 patients with AG had no Btk expression in monocytes (Figure 2) but had mutations in Btk gene. Ten of 11 cases had very low expression of B cells (<1.59%). Five of them had normal Btk expression in monocytes with no mutation in Btk gene. AG cases with Btk gene mutation were diagnosed as X-LA. For the remaining 6 patients, bone marrow aspirate analysis is planned with FCM in order to obtain further data with respect to the type of B cell precursor and also mutation analysis in genes (like BLNK) responsible for B cell maturation.

Keywords: agammaglobulinemia, Bruton's, tyrosine kinase, flow cytometry, immunodeficiency




Figure 1: Btk expression in X-LA case is negative in monocytes

Figure 2: Btk expression in AG case is positive in monocytes

Case	B cell % (lymphocyte gate)	Btk (monocyte gate)	Btk mutation	Diagnosis
AG1	0,00	Neg ative	Positive	X-LA
AG2	0,03	Negative	Positive	X-LA
AG3	11,22	Positive	Negative	
AG4	0,05	Negative	Positive	X-LA
AG5	0,00	Negative	Positive	X-LA
AG6	00,0	Positive	Negative	
AG7	0,05	Positive	Negative	
AG8	1,59	Positive	Negative	
AG9	0,00	Negative	Positive	X-LA
AG10	1,59	Positive	Negative	
AG11	0,48	Positive	Negative	

Table 1: AG cases analysed by FCM

Molecular Immunology & Immunogenetics Congress, 2012

# TURKISH JOURNAL of IMMUNOLOGY

The Official Journal of the Turkish Society of Immunology







Türk İmmünoloji Dergisi – Volume: 1, Number: 17

## TURKISH JOURNAL OF IMMUNOLOGY

# The Official Journal of the Turkish Society of Immunology Volume: 1, Number: 17, Supplement

Molecular Immunology & Immunogenetics Congress – Abstract Book, 2012

Editors-in-Chief: Gunnur DENIZ, PhD, Professor

Managing Editor: H. Barbaros ORAL, MD PhD, Professor

## **Editorial Reviewer Board**

Ahmet GÜL, Turkey
Aydan İKİNCİOĞULLARI, Turkey
Caner SUSAL, Germany
Cezmi AKDİŞ, Switzerland
Dicle GÜÇ, Turkey
Ender TERZİOĞLU, Turkey
Gaye ERTEN, Turkey
Güher SARUHAN DİRESKENELİ, Turkey
Gülderen YANIKKAYA DEMİREL, Turkey
Haner DİRESKENELİ, Turkey
İlhan TEZCAN, Turkey
Jagues PILOT, France
Jon D. LAMAN, the Netherlands
Mahmut ÇARİN, Turkey

Mübeccel AKDİŞ, Switzerland
Necil KÜTÜKÇÜLER, Turkey
Nerin BAHÇECİLER, Republic of Northern Cyprus
Özden SANAL, Turkey
Peter M. Johnson, UK
Pier L. MELONI, Italy
Şebnem KILIÇ, Turkey
Selim Badur, Turkey
Stanimir KYURKCHIEV, Bulgaria
Stefan KAUFMANN, Germany
Stephen E. CHRISTMAS, UK
Şefik Şanal ALKAN, Switzerland
Tevfik DORAK, USA
Yehuda SHOENFELD, Israel